


Pediatric Cardiac Arrest Algorithm for Suspected or Confirmed COVID-19 Patients

Updated April 2020



CPR Quality

- Push hard ($\geq\frac{1}{3}$ of anteroposterior diameter of chest) and fast (100-120/min) and allow complete chest recoil.
- Minimize interruptions in compressions.
- Avoid excessive ventilation.
- Change compressor every 2 minutes, or sooner if fatigued.
- If no advanced airway, 15:2 compression-ventilation ratio.

Shock Energy for Defibrillation

First shock 2 J/kg, second shock 4 J/kg, subsequent shocks ≥ 4 J/kg, maximum 10 J/kg or adult dose

Advanced Airway

- **Minimize closed-circuit disconnection**
- **Use intubator with highest likelihood of first pass success**
- **Consider video laryngoscopy**
- **Prefer cuffed endotracheal tube if available**
- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement
- Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions

Drug Therapy

- **Epinephrine IO/IV dose:** 0.01 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration). Repeat every 3-5 minutes.
- **Amiodarone IO/IV dose:** 5 mg/kg bolus during cardiac arrest. May repeat up to 2 times for refractory VF/pulseless VT.
- **Lidocaine IO/IV dose:** Initial: 1 mg/kg loading dose. Maintenance: 20-50 mcg/kg per minute infusion (repeat bolus dose if infusion initiated >15 minutes after initial bolus therapy).

Return of Spontaneous Circulation (ROSC)

- Pulse and blood pressure
- Spontaneous arterial pressure waves with intra-arterial monitoring

Reversible Causes

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypoglycemia
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary